

## CONTENTS

(Abstracts/contents lists published in: *Chem. Abstr.; Curr. Contents, Phys. Chem. Earth Sci., Life Sci.*)

Introduction . . . . .	IX
An asymmetric approach to 2-deoxynucleosides via organosulfur building blocks as chemical chameleons B. M. Trost and C. Nübling (Stanford, CA, U.S.A.) . . . . .	1
<i>De novo</i> synthesis of carbohydrates by stereoselective aldol reaction: L-cladinose S. H. Montgomery, M. C. Pirring, and C. H. Heathcock (Berkeley, CA, U.S.A.) . . . . .	13
Synthesis of L-gulose, L-galactose, and their acetylated aldehydo forms from 6-S-phenyl-6-thio-D-hexoses F. Santoyo González and H. H. Baer (Ottawa, Ont., Canada) . . . . .	33
Multiple and long-range participation of benzyl groups in intramolecular C-arylation reactions of benzylated glycosides O. R. Martin, S. P. Rao, C. A. V. Hendricks, and R. E. Mahnken (Binghampton, NY, U.S.A.) . . . . .	49
Novel methods for the preparation of partially acetylated carbohydrates S. Hanessian and M. Kagotani (Montreal, Que., Canada) . . . . .	67
1,6-Anhydro- $\beta$ -D-glucopyranose in organic synthesis: preparation of a fragment for the synthesis of rosaramycin G. Procter, D. Genin (Salford, Gt. Britain), and S. Challenger (Cardiff, Gt. Britain) . . . . .	81
Inversion of configurations of contiguous carbinol centers: application to the synthesis of both enantiomers of natural products from the same enantiomerically pure starting material P. Prasit, G. Robertson, and J. Rokach (Pointe Claire-Dorval, Que., Canada) . . . . .	93
Effect on human liver glycosidases and short syntheses of 1a,2a,6a,7a $\beta$ -1,2,6,7-tetrahydroxypyrrolizidine from D-glycero-D-gulo-heptono-1,4-lactone W. F. Collin, G. W. J. Fleet, M. Haraldsson (Oxford, Gt. Britain), I. Cenci di Bello, and B. Winchester (London, Gt. Britain) . . . . .	105
Synthesis of 1-deoxy-6-epicastanospermine and 1-deoxy-6,8a-diepicastanospermine K. H. Aamlid, L. Hough, and A. C. Richardson (London, Gt. Britain) . . . . .	117
Derivatives of methyl $\beta$ -lactosides as substrates for and inhibitors of $\beta$ -D-galactosidase from <i>E. coli</i> K. Bock (Copenhagen, Denmark), and K. Adelhorst (Lyngby, Denmark) . . . . .	131
The use of <i>N</i> -alkoxycarbonyl derivatives of 2-amino-2-deoxy-D-glucose as donors in glycosylation reactions P. Boullanger, M. Jouineau, B. Bouammali, D. Lafont, and G. Descotes (Villeurbanne, France) . . . . .	151
Benzeneselenenyl triflate as an activator of thioglycosides for glycosylation reactions Y. Ito, T. Ogawa (Wako, Japan), M. Numata, and M. Sugimoto (Tokorozawa, Japan) . . . . .	165

Total synthesis of globotriaosylceramide (Gb <sub>3</sub> ) and lysoglobotriaosylceramide (lysoGb <sub>3</sub> ) K. C. Nicolaou, T. J. Caulfield, and H. F. Katoaka (La Jolla, CA, U.S.A.) . . . . .	177
Muramic acid derivatives as glycosyl donors for the synthesis of muramyl-containing glycosphingolipids and fatty acids A. Toepfer and R. R. Schmidt (Konstanz, F.R.G.) . . . . .	193
Synthesis of glycopeptides with the T <sub>N</sub> and T antigen structures, and their coupling to bovine serum albumin H. Kunz, S. Birnbach, and P. Wernig (Mainz, F.R.G.) . . . . .	207
Use of the methylsulfenyl cation as an activator for glycosylation reactions with alkyl (aryl) 1-thioglycopyranosides: synthesis of methyl O-(2-acetamido-2-deoxy- $\beta$ -D-glucopyranosyl)-(1 $\rightarrow$ 6)-O- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 2)- $\alpha$ -D-glucopyranoside, a derivative of the core trisaccharide of <i>E. coli</i> K 12 F. Dasgupta and P. J. Garegg (Stockholm, Sweden) . . . . .	225
1,3,4,6-Tetra-O-acetyl-2-chloroacetamido-2-deoxy- $\beta$ -D-glucopyranose as a glycosyl donor in synthesis of oligosaccharides F. Dasgupta and L. Anderson (Madison, WI, U.S.A.) . . . . .	239
Synthesis of trisaccharide C-D-E and tetrasaccharide B-C-D-E fragments found in orthosomysins M. Trumtel, P. Tavechia, A. Veyrières, and P. Sinay (Paris, France) . . . . .	257
<i>Author index</i> . . . . .	c1
<i>Subject index</i> . . . . .	c3

